

MARKING GUIDELINES

NASIENRIGLYNE

EXAMINATION	NATIONAL SENIOR CERTIFICATE
GRADE / GRAAD	12
DATE / DATUM	JUNE / JUNIE 2024
SUBJECT / VAK	PHYSICAL SCIENCES <i>FISIESE WETENSKAPPE</i>
PAPER / VRAESTEL	2
MARK TOTAL / PUNTE TOTAAL	150
DURATION (HOURS) / DUUR (URE)	3
NUMBER OF PAGES / AANTAL BLADSYE	17



SOUTH AFRICAN COMPREHENSIVE ASSESSMENT INSTITUTE
SUID-AFRIKAANSE KOMPREENSIEWE ASSESSERINGSINSTITUUT

PHYSICAL SCIENCES CHEMISTRY (P2)
FISIESE WETENSKAPPE CHEMIE (V2)


DATE / DATUM: JUNE 2024
QUESTION 1/VRAAG 1

MARKS / PUNTE: 150


- | | | |
|------|-----|-----|
| 1.1 | B✓✓ | (2) |
| 1.2 | A✓✓ | (2) |
| 1.3 | B✓✓ | (2) |
| 1.4 | D✓✓ | (2) |
| 1.5 | A✓✓ | (2) |
| 1.6 | C✓✓ | (2) |
| 1.7 | A✓✓ | (2) |
| 1.8 | D✓✓ | (2) |
| 1.9 | B✓✓ | (2) |
| 1.10 | B✓✓ | (2) |

[20]

QUESTION 2/VRAAG 2

- 2.1 Saturated ✓
 Versadig ✓ (1)
- 2.2.1 Haloalkanes / Alkyl Halides ✓
 Haloalkane/ Alkielhaliede ✓ (1)
- 2.2.2 Carboxyl group ✓
 Karboksiel groep ✓ (1)
- 2.2.3 Ethanol ✓
 Ethanoic acid ✓
 Etanol ✓
 Etanoësuur ✓ (2)
- 2.3.1  ✓
 MARKING GUIDELINES:
 Functional group
 Correct structure (2)
- 2.3.2 $\text{CH}_2(\text{Cl})\text{CH}(\text{CH}_3)\text{CH}_3$ ✓✓
 OF
 $\text{CH}_3\text{CH}(\text{CH}_3)\text{CH}_2\text{Cl}$ ✓✓ (2)
- 2.3.3 $\text{C}_4\text{H}_8\text{O}_2$ ✓✓ (2)
- 2.4 Organic molecules with the same molecular formula, ✓ but different structural formulae. ✓
 Organiese molekule met dieselfde molekulêre formule ✓, maar verskillende struktuur formules ✓ (2)
- 2.5.1 Positional isomers ✓
 Posisie isomere ✓ (1)
- 2.5.2 Functional isomers ✓
 Funksionele isomere ✓ (1)

2.6

**MARKING GUIDELINES:**

- ✓ Functional group
- ✓ Correct structure

(2)

[17]

QUESTION 3/VRAAG 3

- 3.1.1 Boiling point ✓
Kookpunt ✓ (1)
- 3.1.2 The temperature ✓ at which the vapour pressure of a substance equals atmospheric pressure. ✓
Die temperatuur ✓ waar die dampdruk van die stof gelyk is aan atmosferiese druk ✓ (2)
- 3.1.3 A ✓ (1)
- 3.1.4 Pent-2-ene is a non-polar alkene.
 It has weak London forces between its molecules. ✓
Pentan-2-ol has stronger ✓ hydrogen bonds ✓ between its molecules.
More energy needed to overcome the forces in pentan-2-ol. ✓
*Pent-2-een is 'n nie-polêre alkeen
 Daar is swak London kragte tussen die molekule ✓
Pentan-2-ol besit sterker ✓ waterstofbindings ✓ tussen die molekule
Meer energie word benodig om intermolekulêre kragte in pentan-2-ol te oorkom ✓* (4)
- 3.2.1 2,2,4-trimethylpentane ✓ ✓
2,2,4-trimetielpentaaan ✓ ✓ (2)
- 3.2.2 Chain isomer ✓
Ketting isomeer ✓ (1)
- 3.3.1 The pressure exerted by a vapour at equilibrium with its liquid in a closed system. ✓ ✓
Die druk wat 'n ingeslote damp wat in ewewig met sy vloeistof is, uitoefen op die oppervlak van die vloeistof ✓ ✓ (2)
- 3.3.2 Pentan-1-ol ✓
Pentan-1-ol ✓ (1)

3.3.3 The predominant intermolecular forces in 2,2-dimethylbutane are weak London forces. ✓

Ethyl propanoate has dipole-dipole ✓ forces between its molecules.

More energy is needed to overcome the dipole-dipole forces ✓ in ethyl propanoate.

Therefore ethyl propanoate has a higher boiling point and a lower vapour pressure ✓ than 2,2-dimethylbutane.

2,2-dimetiëlbutaan besit swak London kragte ✓

Etiëlpropanoaat besit sterker dipool-dipool kragte ✓

Meer energie word benodig om die dipool-dipool kragte ✓ in etiëlpropanoaat te oorkom

Etiëlpropanoaat besit hoër kookpunt en dus 'n laer dampdruk as ✓

2,2-dimetiëlbutaan

(4)

[18]

QUESTION 5/VRAAG 5

5.1 $n = \frac{m}{M}$ ✓
 $= \frac{12}{100}$ ✓
 $= 0,12 \text{ mol}$ ✓ (3)

5.2 $n(\text{CaCO}_3) = 0,12 \text{ mol}$
 $n(\text{HCl}) = c \times V$
 $= 3 \times 0,2$ ✓
 $= 0,6 \text{ mol}$ ✓
 $\text{CaCO}_3 : \text{HCl}$
 $1 : 2$
 $0,12 : 0,24$ ✓
 $0,24 < 0,6$ HCl is in excess/ oormaat ✓ (4)

5.3 4 minutes has a higher rate ✓ than 8 minutes
4 minute het 'n hoër reaksietempo ✓ as 8 minute (1)

5.4 As the reaction proceeds, the concentration of the HCl decreases. ✓
 This means fewer particles per volume. ✓
Fewer effective collisions per unit time. ✓
 Which results in a lower rate of reaction at 8 minutes
Soos reaksie verloop, neem die HCl konsentrasie af ✓
Dit beteken minder deeltjies per volume ✓
Minder effektiewe botsings per tyds eenheid ✓
Dus 'n laer reaksie tempo by 8 minute (3)

5.5 3,25 g ✓ (1)

5.6 $n_{(\text{CaCO}_3)} = 0,12 \text{ mol}$
 $n_{(\text{CO}_2)} = n_{(\text{CaCO}_3)} = 0,12 \text{ mol} \checkmark$
 $m_{(\text{CO}_2)} = n \times M$
 $= 0,12 \times 44 \checkmark$
 $= 5,28 \text{ g}$

$$\% \text{ yield} = \frac{\text{actual mass}}{\text{theoretical mass}} \times 100 \checkmark \quad (5)$$

$$= \frac{3,25}{5,28} \times 100 \checkmark$$

$$= 61,55 \% \checkmark$$

5.7.1 Steeper \checkmark

Steiler/ hoër gradient \checkmark (1)

5.7.2 No change \checkmark

Geen verandering \checkmark (1)

[19]

QUESTION 6/VRAAG 6

6.1.1 A reaction is reversible when products can be converted ✓ back to reactants and vice versa. ✓

'n Reaksie is omkeerbaar wanneer die produkte terug na reaktante omgeskakel kan word ✓ en vice versa. ✓ (2)

6.1.2 $2 \text{NO} (\text{g}) \rightarrow \text{N}_2 (\text{g}) + \text{O}_2 (\text{g})$ ✓ (1)

6.1.3 The system is in (dynamic) chemical equilibrium. ✓
The rate of reverse and forward reactions is equal. ✓

*Die sisteem is in (dinamiese) chemiese ewewig ✓
Die tempo van die voorwaartse en terugwaartse reaksie is dieselfde ✓* (2)

6.1.4 Decreased. ✓

Afneem ✓ (1)

6.1.5 Exothermic. ✓

Eksotermies ✓ (1)

6.1.6 Rates of both forward and reverse reactions decrease.
Reverse reaction rate decreases more. ✓
Rate of endothermic reaction will decrease more to absorb the energy.
Thus, the reverse reaction is endothermic. ✓
Forward reaction was (increased) favoured by the decreases in temperature.

*Die tempo van beide die voorwaartse en terugwaartse reaksies neem af
Die terugwaartse reaksie tempo neem meer af ✓
Die tempo van die endotermiese reaksie sal meer afneem omdat die energie te absorber (Volgens Le Chatelier se beginsel sal temperatuur verhoging die endotermiese reaksie bevoordeel)
Dus die terugwaartse reaksie is endotermies ✓
Die voorwaartse reaksie is bevoordeel deur die afname in temperatuur* (2)

6.1.7 K_c will increase. ✓

Since $K_c = \frac{[NO]^2}{[N_2][O_2]}$ and the forward reaction is favoured, it will cause an increase in $[NO]$ ✓, value of K_c will increase.

OR

Decrease in the $[N_2]$ and $[O_2]$ ✓, value of K_c will increase.

K_c sal toeneem ✓

Omdat $K_c = \frac{[NO]^2}{[N_2][O_2]}$ is en die voorwaatse reaksie is bevoordeel sal $[NO]$ toeneem en K_c sal toeneem ✓

OF

Afname in die $[N_2]$ en $[O_2]$ en sal K_c sal toeneem ✓ (2)

6.1.8 Catalyst was added/ Pressure is increased. ✓

Katalisator is bygevoeg / druk word verhoog ✓ (1)

6.1.9 The rates of both the forward and reverse reactions increased ✓✓

Die tempo van beide die voorwaartse en terugwaartse reaksie verhoog ✓✓ (2)

6.2.1

	NO₂	NO	N₂O	O₂
Ratio	1	1	1	1
Initial concentration / Oorspronklike konsentrasie (mol·dm ⁻³)	0,06	0,29	0,18	0,38
Change in concentrations/ Verandering in konsentrasie	+ 0,06	+ 0,06	- 0,06	- 0,06
Equilibrium concentration / Konsentrasie by ewewig	0,12	0,35	0,12	0,32

$$K_c = \frac{[N_2O][O_2]}{[NO_2][NO]}$$

$$= \frac{(0,12)(0,32)}{(0,12)(0,35)}$$

$$= 0,91$$

MARKING GUIDELINES:

- ✓ Use of ratio
- ✓ Concentrations of NO, N₂O and O₂ at equilibrium
- ✓ K_c expression
- ✓ Substitution of values
- ✓ Answer (no unit)

(5)

6.2.2 Reverse reaction ✓

Terugwaartse reaksie ✓

(1)

6.2.3 $[\text{NO}_2]$ increased from $0,06 \text{ mol}\cdot\text{dm}^{-3}$ to $0,12 \text{ mol}\cdot\text{dm}^{-3}$. ✓
 NO_2 is the product of the reverse reaction.

OR

K_c value decreased. ✓

*$[\text{NO}_2]$ neem toe van $0,06 \text{ mol}\cdot\text{dm}^{-3}$ na $0,12 \text{ mol}\cdot\text{dm}^{-3}$ ✓
 NO_2 word deur die terugwaartse reaksie gevorm*

OF

K_c waarde het afgeneem ✓

(1)

[21]

QUESTION 7/VRAAG 7

7.1.1 An acid is a proton donor. ✓ ✓

'n Suur is 'n proton skenker ✓✓ (2)

7.1.2 H_3PO_4 can donate more than one proton (H^+ -ion) per molecule. ✓ H_3PO_4 kan meer as een proton (H^+ -ioon) per molekule skenk ✓ (1)

7.2.1
$$c = \frac{m}{MV} \checkmark$$

$$0,22 = \frac{m}{40 \times 0,5} \checkmark$$

$$m = 4,4 \text{ g} \checkmark \quad (3)$$

7.2.2
$$\frac{n_a}{n_b} = \frac{c_a V_a}{c_b V_b} \checkmark$$

$$\frac{1}{3} \checkmark = \frac{c_a \times 20}{0,22 \times 15} \checkmark \checkmark$$

$$c_a = 0,055 \text{ mol} \cdot \text{dm}^{-3} \checkmark$$

OR

$$n_b = c \cdot V \checkmark$$

$$= 0,22 \times 0,015 \checkmark$$

$$= 3,3 \times 10^{-3} (0,0033) \text{ mol}$$

$$n_b : n_a$$

$$3 : 1 \checkmark$$

$$3,3 \times 10^{-3} : 1,1 \times 10^{-3} (0,0011)$$

$$c_a = \frac{n}{V}$$

$$= \frac{1,1 \times 10^{-3}}{0,020} \checkmark$$

$$= 0,055 \text{ mol} \cdot \text{dm}^{-3} \checkmark$$

MARKING GUIDELINES:

- ✓ Formula
- ✓ Ratio
- ✓✓ Substitution
- ✓ Answer with unit

(5)

MARKING GUIDELINES:

- ✓ Any Formula
- ✓ substitution
- ✓ Ratio
- ✓ Substitution
- ✓ Answer with unit

(5)

7.2.3 Greater than 7. ✓

Groter as 7 ✓

(1)

- 7.2.4 NaOH is a strong base and H₃PO₄ is a weak acid. ✓
PO₄³⁻ will react with water to produce OH⁻¹ ions.

$$\text{PO}_4^{3-} + 3\text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{PO}_4 + 3\text{OH}^-$$
 ✓
Excess OH⁻ in solution results in a basic solution. ✓
 Indicator that changes colour in basic range (phenolphthalein) required.

NaOH is 'n sterk basis en H₃PO₄ is 'n swak suur ✓
PO₄³⁻ sal met water reageer om OH⁻¹-ione te vorm

$$\text{PO}_4^{3-} + 3\text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{PO}_4 + 3\text{OH}^-$$
 ✓
Dit is 'n basiese oplossing omdat daar 'n oormaat OH⁻¹-ione in die oplossing is ✓
'n Indikator wat kleur verander in die pH- gebied(basiese gebied) word benodig (3)

[15]


QUESTION 8/VRAAG 8

- 8.1 Chemical energy is converted in electrical energy. ✓ ✓
Chemiese energie word omgeskakel in elektriese energie. ✓ ✓ (2)
- 8.2 1 mol·dm⁻³ ✓ (1)
- 8.3 A substance of which the aqueous solution contains ions. ✓ ✓
OR
 A substance that dissolves in water to give a solution that conducts electricity. ✓ ✓
'n Stof wat in 'n waterige oplossing ione besit ✓ ✓
OR
'n Stof wat oplosbaar is in water en 'n oplossing vorm wat elektrisiteit gelej ✓ ✓ (2)
- 8.4 Co / cobalt ✓
 Co / kobalt ✓ (1)
- 8.5 $\text{Cl}_2 + 2\text{e}^- \rightarrow 2\text{Cl}^-$ ✓ ✓ (2)
- 8.6 $\text{Co (s)}|\text{Co}^{2+} (\text{aq}) || \text{Cl}_2 (\text{g}) | \text{Cl}^- (\text{aq})$, Pt (s) ✓ at 25 °C
- MARKING GUIDELINES:**

 - ✓ Anode up to saltbridge
 - ✓ Cathode
 - ✓ Pt
- (3)
- 8.7 $E^{\theta}_{\text{cell}} = E^{\theta}_{\text{cathode}} - E^{\theta}_{\text{anode}}$ ✓
 $= 1,36 \text{ ✓} - (-0,28) \text{ ✓}$
 $= 1,64 \text{ V ✓}$
- MARKING GUIDELINES:**


 - ✓ Formula
 - ✓ Substitution
 - ✓ Answer with unit
- (4)

8.8 Suggestion is not correct ✓



Increasing $[\text{Co}^{2+}]$ will favour the reverse reaction ✓ decreasing the emf. ✓

Die voorstel is nie korrek nie ✓



'n Toename in die $[\text{Co}^{2+}]$ sal die terugwaartse reaksie bevoordeel ✓ en die emk sal afneem ✓

(3)

[18]

QUESTION 9/VRAAG 9

9.1 Chromium (Cr)✓

Chroom (Cr) ✓

(1)

9.2 Cr³⁺ is a much stronger oxidising agent than water. ✓
Cr³⁺ will be reduced at the cathode/key✓*Cr³⁺ is 'n sterker oksideermiddel as water ✓**Cr³⁺ sal gereduseer word by die katode/sleutel ✓*

(3)

9.3 Cr: electrons/ elektrone

1 :3

0,073 : 0.22✓

$$M(\text{Cr}) = n \times M \checkmark$$

$$= 0,073 \times 52 \checkmark$$

$$= 3,796 \text{ g} \checkmark$$

(4)

MARKING GUIDELINES:

- ✓ Use of ratio
- ✓ Formula
- ✓ Substitution
- ✓ Answer with unit

[8]