


NASIENRIGLYNE

EKSAMEN		NATIONALE SENIOR SERTIFIKAAT	
GRAAD		12	
DATUM		JUNIE 2024	
VAK		WISKUNDE	
VRAESTEL		1	
PUNTE TOTAAL		150	
TYD (URE)		3	
AANTAL BLADSYE		11	



SOUTH AFRICAN COMPREHENSIVE ASSESSMENT INSTITUTE
SUID-AFRIKAANSE KOMPREENSIEWE ASSESSERINGSINSTITUUT

VRAAG 1

1.1.1	$(x - 5)(x + 2) = 0$ $x = 5$ or $x = -2$	✓ Elke faktor = 0 ✓ Antwoorde (2)
1.1.2	Stel $\sqrt{p} = x$ $\sqrt{p} = 5$ or $\sqrt{p} = -2$ $p = 25$ geen antwoord	✓ Stel p ✓ ✓ Antwoord (3)
1.2.1	$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $x = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(2)(-8)}}{2(2)}$ $x = \frac{3 \pm \sqrt{73}}{4}$ $x = 2,89$ or $x = -1,39$	✓ Vervang in formule ✓ Vereenvoudig ✓ Antwoorde (3)
1.2.2	$x^2 + 2x - 80 \geq 0$ $(x - 8)(x + 10) \geq 0$ $x \leq -10$ or $x \geq 8$ 	✓ Faktoreer ✓ 1 vir waardes ✓ 1 vir notasie (3)
1.3	$y = 5 - 2x$ $4x^2 - 2(5 - 2x)^2 = 46$ $4x^2 - 50 + 40x - 8x^2 = 46$ $x^2 - 10x + 24 = 0$ $(x - 4)(x - 6) = 0$ $x = 4$ or/ of $x = 6$ $y = -3$ or/ of $y = -7$ OF $x = \frac{5}{2} - \frac{y}{2}$ $4\left(\frac{5}{2} - \frac{y}{2}\right)^2 - 2y^2 = 46$ $25 - 10y + y^2 - 2y^2 = 46$ $y^2 + 10y + 21 = 0$ $(y + 3)(y + 7) = 0$ $y = -3$ or $y = -7$ $x = 4$ or $x = 6$ $b = \frac{1}{2}$ or $b = -3$ and $a = \frac{9}{4}$ or $a = 4$	✓ $x = 5 - 2y$ ✓ Vervang ✓ Standaard vorm ✓ Faktoreer ✓ x -waardes ✓ y waardes (6)

1.4	$\sqrt{\frac{25 \cdot 5^{1003} + 625 \cdot 5^{1003}}{26(5^{1003})}}$ $= \sqrt{\frac{5^{1003}(25+625)}{26(5^{1003})}}$ $= \sqrt{\frac{650}{26}}$ $= 5$	<ul style="list-style-type: none"> ✓ Eksponent 5^{1002} ✓ Faktoriseer ✓ 650 ✓ Antwoord (4)
1.5	$\frac{b}{a} = \frac{c}{b}$ $\therefore b^2 = ac$ $\Delta = b^2 - 4ac = ac - 4ac$ $= -3ac$ <p>Omdat a en c negatief is, is Δ negatief. Dus geen reële wortels nie.</p>	<ul style="list-style-type: none"> ✓ Meetkundige ry ✓ b^2 ✓ Δ ✓ Gevolgtrekking (4)
[25]		

VRAAG 2

2.1	$4x - x - 2 = 6x + 4 - 4x$ $x = 6$ <p>8; 24; 40; ...</p>	✓ Vergelyking ✓ x ✓ Ry (3)
2.2	$a = \frac{5}{3} \text{ and } r = 3$ $S_n = \frac{a(r^n - 1)}{r - 1} = \frac{\frac{5}{3}(3^n - 1)}{2} = \frac{1820}{3}$ $r^n = 729 = 3^6$ $n = 6$	✓ ✓ a en r ✓ Vervang ✓ $= \frac{1820}{3}$ ✓ 729 ✓ Antwoord (6)
2.3.1	$-1 < \frac{2x - 5}{2} < 1$ $-2 < 2x - 5 < 2$ $\frac{3}{2} < x < \frac{7}{2}$	✓ Beperkings ✓ Maal 2 ✓ Antwoord (3)
2.3.2	$S_\infty = \frac{1}{1 - \frac{2x - 5}{2}} = \frac{4}{9}$ $\frac{2}{7 - 2x} = \frac{4}{9}$ $18 = 28 - 8x$ $x = \frac{5}{4}$	✓ Formule ✓ ✓ Vereenvoudig ✓ Antwoord (4)
[16]		


VRAAG 3

3.1	$d + k + p; 4d + 2k + p; 9d + 3k + p$ $3d + k; 5d + k$ $\swarrow \nearrow$ $2d$	✓ ✓ eerste 3 terme ✓ ✓ 1 st verskil (4)
3.2.1	Louisa: Meetkundig. $27 \times 3 = 81$ Thabo: Kwadraties. $3 + 6 = 9; 9 + 18 = 27; 27 + 30 = 57$	✓ Louisa ✓ Thabo (2)
3.2.2	$3; 9; 27; 57.....$ $\swarrow \nearrow \swarrow \nearrow \swarrow \nearrow$ $6 \quad 18 \quad 30$ $\swarrow \nearrow \swarrow \nearrow$ $12 \quad 12$ $2a = 12, a = 6$ $3a + b = 6, b = -12$ $a + b + c = 3, c = 9$ $T_{10} = 6(10)^2 - 12(10) + 9$ $= 489$	✓ ✓ ✓ a, b en c ✓ $n = 10$ ✓ Antwoord (4)
[10]		

VRAAG 4

4.1.1	$A = 940000 \left(1 - \frac{d}{100}\right)^7 \text{ ou}$ $A = 940000(1 + 0,12)^7 = R2078040,52 \text{ nuwe}$ $R2078040,52 - 940000 \left(1 - \frac{d}{100}\right)^7 = 1843712,18$ $\left(1 - \frac{d}{100}\right)^7 = 0,2492 \dots$ $\frac{d}{100} = 0,18000000$ $d = 18\%$	<ul style="list-style-type: none"> ✓ Ou ✓ Nuwe ✓ Vergelyking ✓ Vereenvoudig ✓ Antwoord (5)
4.1.2	$1843712,18 = x \cdot \frac{1,0075^{84} - 1}{0,0075}$ $x = R15835,79$	<ul style="list-style-type: none"> ✓ Regte vervanging in regte formule ✓ Antwoord (2)
4.2.1	$22000(1,0125) = x \cdot \frac{1 - (1 + 0,0125)^{-47}}{0,0125}$ $x = R629,58$ <p>Totaal = R629,58 x 47 = R29590,35</p>	<ul style="list-style-type: none"> ✓ Regte r ✓ Regte formule met korrekte vervanging ✓ antwoord x ✓ x47 antwoord (4)
4.2.2	$OB = 22000(1,0125)^{24} - 629,58 \frac{1,0125^{23} - 1}{0,0125}$ $= R29641,72 - R16657,03$ $= R12984,69$ <p>Nee, hy sal nie kan nie</p>	<ul style="list-style-type: none"> ✓ Waarde van rekenaar ✓ Bedrag betaal ✓ Berekeninge ✓ Antwoord ✓ Nee (5)
[16]		

VRAAG 5

5.1.3	$y = \frac{a}{x+2} + 4$ $2 = \frac{a}{0+2} + 4$ $\frac{a}{2} = -2$ $a = -4; p = 2; q = 4$	✓ ✓ p en q ✓ Vervang (0; 2) ✓ $a = -4$ (4)
5.1.2	$y = x + c; 4 = -2 + c$ $c = 6$	✓ Vervanging ✓ Antwoord (2)
5.2.1	$g(x) = \left(\frac{2}{3}\right)^x$	✓ Antwoord (1)
5.2.2	$f^{-1}(x) = \log_{\frac{3}{2}} x$	✓ ✓ Antwoord (2)
5.2.3		✓ ✓ ✓ 1 vir elke grafiek (3)
[12]		

VRAAG 6

6.1	$f(x) = -(x^2 - 4x + 4) + 4 + 9$ $f(x) = -(x - 2)^2 + 9$	✓ Minus en die 4 ✓ Antwoord (3)
6.2	(2; 9)	✓ ✓ Antwoord (2)
6.3	$x^2 - 4x - 5 = 0$ $(x - 5)(x + 1) = 0$ $x = 5$ or $x = -1$ $-x^3 + 10x^2 - 17x - 28 = -(x + 1)(x^2 - 11x + 28)$ $(x + 1)(x - 4)(x - 7) = 0$ $x = -1$ or $x = 4$ or $x = 7$	✓ Faktore ✓ Antwoorde ✓ Besef $(x + 1)$ is 'n faktor. ✓ ✓ Die ander faktor ✓ Faktoriseer ✓ Antwoorde (7)
6.4	$g'(x) = -3x^2 + 20x - 17 = 0$ $(3x - 17)(x - 1) = 0$ $x = \frac{17}{3}$ or $x = 1$	✓ $f' = 0$ ✓ Faktoriseer ✓ Antwoord (3)
6.5	$h(x) = -(x - 4)^2 + 13$	✓ ✓ Elke waarde (2)
6.6	$4 < x < 5$ or $x > 7$	✓ ✓ ✓ Elke antwoord (3)
6.7	$x = \frac{17}{3}$ Maks draaipunt. Dus $g''(x) < 0$, $g'(x) = 0$ en $g(x) > 0$	✓ Waarde ✓ ✓ Motivering (3)
[23]		

VRAAG 7


<p>7.1</p>	$f(x) = 3x^2 + 1$ $f(x + h) = 3(x + h)^2 + 1$ $= 3x^2 + 6xh + 3h^2 + 1$ $f'(x) = \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h}$ $f'(x) = \lim_{h \rightarrow 0} \frac{3x^2 + 6xh + 3h^2 + 1 - 3x^2 - 1}{h}$ $f'(x) = \lim_{h \rightarrow 0} \frac{6xh + 3h^2}{h}$ $f'(x) = \lim_{h \rightarrow 0} \frac{h(6x + 3h)}{h}$ $f'(x) = \lim_{h \rightarrow 0} (6x + 3h)$ $f'(x) = 6x$	<p>✓ $f(x + h)$</p> <p>✓ regte vervanging</p> <p>✓ Vereenvoudig</p> <p>✓ $(2ax + ah)$</p> <p>✓ $2ax$</p> <p>NB: Penaliseer een keer vir verkeerde of geen notasie.</p> <p>$\left[\lim_{h \rightarrow 0} (\) \right]$ (5)</p>
<p>7.2.1</p>	$f(x) = 2x - \sqrt{x}$ $f(x) = 2x - x^{\frac{1}{2}}$ $f'(x) = 2 - \frac{1}{2}x^{-\frac{1}{2}}$	<p>✓ $x^{\frac{1}{2}}$</p> <p>✓ 2 ✓ $-\frac{1}{2}x^{-\frac{1}{2}}$ (3)</p>
<p>7.2.2</p>	$D_x \left[\frac{x^2 - 3x + 2}{x - 1} \right]$ $D_x \left[\frac{(x - 2)(x - 1)}{x - 1} \right]$ $= 1$	<p>✓ ✓ Faktore</p> <p>✓ Antwoord (3)</p>
<p>7.4.1</p>	<p>$x = -1$ because $f'(-1) = 0$ and $f''(x) < 0$</p>	<p>✓ Waarde</p> <p>✓ ✓ Rede (3)</p>
<p>7.4.2</p>	<p>$x = 2$ because $f''(x) = 0$</p>	<p>✓ ✓ Waarde en 2e afgeleide = 0 (2)</p>

7.4.3	$y = a(x + 1)(x - 5)$ $-9 = a(2 + 1)(2 - 5)$ $a = 1$ $f'(x) = x^2 - 4x - 5$ <p style="text-align: center;">OR</p> $y = a(x - 2)^2 - 9$ $0 = a(5 - 2)^2 - 9$ $a = 1$ $f'(x) = x^2 - 4x - 5$ <p style="text-align: center;">OR</p> $y = a(x - 2)^2 - 9$ $0 = a(-1 - 2)^2 - 9$ $a = 1$ $f'(x) = x^2 - 4x - 5$	<p>✓ 1 Korrekte waardes in korrekte formule</p> <p>✓ (5;0) Vervanging</p> <p>✓ Standaard vorm (3)</p>
7.4.4	$f'(x) = 3ax^2 + 2bx + c$ $a = \frac{1}{3}, b = -2, c = -5$	<p>✓ afgeleide</p> <p>✓ a</p> <p>✓ ✓ b en c (4)</p>
[28]		

VRAAG 8

8.1	$CD = x^2 + 4 - (-x^2 + ax - 5) = 2x^2 - ax + 9$ $\frac{d}{dx}(CD) = 4x - a = 0$ $a = 4x$ <p>En $CD = 7$</p> $2x^2 - 4x^2 + 9 = 7$ $x^2 = 1$ $x = 1$ $a = 4$	<ul style="list-style-type: none"> ✓ CD ✓ Differensieer $CD=0$ ✓ $a = 4x$ ✓ Stel $CD = 7$ ✓ Vervang a ✓ $x = 1$ ✓ $a = 4$ <p style="text-align: right;">(7)</p>
[7]		

VRAAG 9

9.1.1		<ul style="list-style-type: none"> ✓ 9 ✓ 11, 10 en 14 ✓ 5, 6 en 10 ✓ 15 <p style="text-align: right;">(5)</p>
9.1.2	15	<ul style="list-style-type: none"> ✓ Antwoord (1)
9.1.3	<p>21 mense</p> $\therefore \frac{21}{80}$	<ul style="list-style-type: none"> ✓ 21 ✓ Antwoord (2)
9.2.1	10!	<ul style="list-style-type: none"> ✓ Antwoord (1)
9.2.2	<p>Aantal maniere: $(2! \times 3! \times 5!) \times 3!$</p> $= 8640$ $P = \frac{8640}{3628800} = \frac{1}{420}$	<ul style="list-style-type: none"> ✓ ✓ Aantal maniere ✓ 8640 ✓ Waarskynlikheid (4)
[13]		